

APPLICATIONS OF HOFER'S GEOMETRY TO HAMILTONIAN DYNAMICS

URS FRAUENFELDER AND FELIX SCHLENK

ABSTRACT. We prove the following three results in Hamiltonian dynamics.

- The Weinstein conjecture holds true for every displaceable hypersurface of contact type.
- Every magnetic flow on a closed Riemannian manifold has contractible closed orbits for a dense set of small energies.
- Every closed Lagrangian submanifold whose fundamental group injects and which admits a Riemannian metric without closed geodesics has the intersection property.

The proofs all rely on the following creation mechanism for closed orbits: If the ray $\{\varphi_H^t\}$, $t \geq 0$, of Hamiltonian diffeomorphisms generated by a sufficiently nice compactly supported time-independent Hamiltonian stops to be a minimal geodesic in its homotopy class, then a non-constant contractible closed orbit must appear.

1. RESULTS

We consider an arbitrary smooth symplectic manifold (M, ω) . A *hypersurface* S in M is a smooth compact connected orientable codimension 1 submanifold of $M \setminus \partial M$ without boundary. A closed characteristic on S is an embedded circle in S all of whose tangent lines belong to the distinguished line bundle

$$\mathcal{L}_S = \{(x, \xi) \in TS \mid \omega(\xi, \eta) = 0 \text{ for all } \eta \in T_x S\}.$$

Examples show that $\mathcal{L}_S$ might not carry any closed characteristic, see [9, 10]. We therefore follow [15] and consider parametrized neighbourhoods of S . Since S is orientable, there exists an open neighbourhood I of 0 and a smooth diffeomorphism

$$\psi: S \times I \rightarrow U \subset M$$

such that $\psi(x, 0) = x$ for $x \in S$. We call ψ a *thickening* of S , and we abbreviate $S_\epsilon = \psi(S \times \{\epsilon\})$. We denote by $\mathcal{P}^\circ(S_\epsilon)$ the set of closed characteristics on S_ϵ which are contractible in M . Let $\mathcal{H}_c(M)$ be the

Date: February 1, 2008.

set of smooth functions $[0, 1] \times M \rightarrow \mathbb{R}$ whose support is compact and contained in $[0, 1] \times (M \setminus \partial M)$. The Hamiltonian vector field of $H \in \mathcal{H}_c(M)$ defined by

$$\omega(X_{H_t}, \cdot) = -dH_t(\cdot)$$

generates a flow φ_H^t . The set of time-1-maps φ_H form the group

$$\text{Ham}_c(M, \omega) := \{\varphi_H \mid H \in \mathcal{H}_c(M)\}$$

of compactly supported Hamiltonian diffeomorphisms of (M, ω) . We call a subset A of (M, ω) *displaceable* if there exists $\varphi \in \text{Ham}_c(M, \omega)$ such that $\varphi(A) \cap A = \emptyset$. Our basic result is

Theorem 1.1. *Assume that S is a displaceable hypersurface in a symplectic manifold (M, ω) . Then for any displaceable thickening $\psi: S \times I \rightarrow U \subset M$ the set $\{\epsilon \in I \mid \mathcal{P}^\circ(S_\epsilon) \neq \emptyset\}$ is dense in I .*

Theorem 1.1 should be compared with the Hofer–Zehnder Theorem [16, 17, 29] stating that for any thickening $\psi: S \times I \rightarrow U \subset M$ for which the Hofer–Zehnder capacity $c_{\text{HZ}}(U)$ is finite the hypersurface S_ϵ carries a closed characteristic for almost all $\epsilon \in I$. We shall draw three applications from Theorem 1.1.

1. The Weinstein conjecture

A hypersurface S in (M, ω) is called *of contact type* if there exists a Liouville vector field X (i.e., $\mathcal{L}_X \omega = d\iota_X \omega = \omega$) which is defined in a neighbourhood of S and is everywhere transverse to S . Weinstein conjectured in [39] that every hypersurface S of contact type with $H^1(S; \mathbb{R}) = 0$ carries a closed characteristic.

Corollary 1.2. *Assume that $S \subset (M, \omega)$ is a displaceable hypersurface of contact type. Then $\mathcal{P}^\circ(S) \neq \emptyset$. In particular, the Weinstein conjecture holds true for every displaceable hypersurface of contact type.*

Proofs of the Weinstein conjecture for *all* hypersurfaces of contact type of *special classes* of symplectic manifolds have been found in [36, 15, 13, 6, 14, 18, 26, 37, 24, 38, 2, 23, 25]. Corollary 1.2 generalizes the results in [36, 15, 6, 38, 23], where the ambient symplectic manifold is of the form $(M \times \mathbb{R}^2, \omega \oplus \omega_0)$. Under the additional assumption that (M, ω) satisfies $[\omega]|_{\pi_2(M)} = 0$ and is convex, Corollary 1.2 has been proved in [7].

2. Closed trajectories of a charge in a magnetic field

Consider a Riemannian manifold (N, g) of dimension at least 2, and let $\omega_0 = \sum_i dp_i \wedge dq_i$ be the standard symplectic form on the cotangent

bundle T^*N . The motion of a unit charge on (N, g) subject to a magnetic field can be described as the Hamiltonian flow of the Hamiltonian system

$$(1) \quad H: (T^*N, \omega_\sigma) \rightarrow \mathbb{R}, \quad H(q, p) \mapsto \frac{1}{2} |p|^2,$$

where σ is the closed 2-form on N representing the magnetic field and where ω_σ is the twisted symplectic form $\omega_\sigma = \omega_0 + \pi^*\sigma$. A trajectory of a charge on (N, g) in the magnetic field σ has constant speed, and closed trajectories γ on N of speed $c > 0$ correspond to closed orbits of (1) on the energy level $E_c = \{H = c^2/2\}$. An old problem in Hamiltonian mechanics asks for closed orbits on a given energy level E_c , see [8]. We denote by $\mathcal{P}^\circ(E_c)$ the set of closed orbits on E_c which are contractible in T^*N and hence project to contractible closed trajectories on N .

Corollary 1.3. *Consider a closed Riemannian manifold (N, g) endowed with a closed 2-form σ which does not vanish identically. There exists $d > 0$ such that $\mathcal{P}^\circ(E_c) \neq \emptyset$ for a dense set of $c \in]0, d[$.*

The number $d > 0$ has a geometric meaning: If the Euler characteristic $\chi(N)$ vanishes, d is the supremum of the real numbers c for which the sublevel set

$$H^c = \{(q, p) \in T^*N \mid H(q, p) = \frac{1}{2}|p|^2 \leq c\}$$

is displaceable in (T^*N, ω_σ) , and if $\chi(N)$ does not vanish, d is defined via stabilizing (1) by $(T^*S^1, dy \wedge dx) \rightarrow \mathbb{R}, (x, y) \mapsto \frac{1}{2}|y|^2$. Corollary 1.3 generalizes various other results on closed orbits of magnetic flows on small energy levels: The existence of a sequence $c \rightarrow 0$ with $\mathcal{P}^\circ(E_c) \neq \emptyset$ has been proved by Polterovich [34] and Macarini [27] under the assumption that $[\sigma]|_{\pi_2(M)} = 0$ and by Ginzburg–Kerman [11] under the assumption that σ is symplectic. If σ is symplectic and $[\sigma]|_{\pi_2(M)} = 0$, Corollary 1.3 has been proved in [3]. For exact forms [7] and rational symplectic forms [10, 27, 28] it is known that $\mathcal{P}^\circ(E_c) \neq \emptyset$ for almost all sufficiently small $c > 0$, and if σ is symplectic and compatible with g , then $\mathcal{P}^\circ(E_c) \neq \emptyset$ for all sufficiently small $c > 0$, see [19]. We refer to [8] and [7, Section 12.4] for more details on the state of the art of the existence problem for closed trajectories of a charge in a magnetic field.

3. Lagrangian intersections

A middle-dimensional submanifold L of (M, ω) is called *Lagrangian* if ω vanishes on L . According to a celebrated theorem of Gromov, [12, 2.3.B'_3], a closed Lagrangian submanifold $L \subset M \setminus \partial M$ with

$[\omega]|_{\pi_2(M,L)} = 0$ is not displaceable. The following result generalizes Theorem 1.4.A in [21] and Theorem 13.1 in [7].

Corollary 1.4. *Assume that $L \subset M \setminus \partial M$ is a closed Lagrangian submanifold of (M, ω) such that*

- (i) *the injection $L \subset M$ induces an injection $\pi_1(L) \subset \pi_1(M)$;*
- (ii) *L admits a Riemannian metric none of whose closed geodesics is contractible.*

Then L is not displaceable.

Acknowledgements. We cordially thank Leonid Polterovich for pointing out to us that the use of functions as in the figure below in his approach to closed orbits of a charge in a magnetic field will provide a result like Corollary 1.3. This paper was written during the second authors stay at Tel Aviv University in April 2003. He wishes to thank Hari and Harald and TAU for their warm hospitality.

2. PROOFS


Proof of Theorem 1.1: We closely follow Polterovich's beautiful argument in [34, Section 9.A]. We fix $\epsilon_0 \in I$ and choose $\delta > 0$ so small that $[\epsilon_0 - 2\delta, \epsilon_0 + 2\delta] \subset I$. Let $f: \mathbb{R} \rightarrow [0, 1]$ be a smooth function such that

$$f(\epsilon) = 0 \text{ if } \epsilon \notin]\epsilon_0 - 2\delta, \epsilon_0 + 2\delta[, \quad f(\epsilon) = 1 \text{ if } \epsilon \in [\epsilon_0 - \delta, \epsilon_0 + \delta],$$

and

$$f'(\epsilon) > 0 \text{ if } \epsilon \in]\epsilon_0 - 2\delta, \epsilon_0 - \delta[, \quad f'(\epsilon) < 0, \text{ if } \epsilon \in]\epsilon_0 + \delta, \epsilon_0 + 2\delta[,$$

see the figure below.



We define the time-independent Hamiltonian $F \in \mathcal{H}_c(M)$ by

$$F(x) = \begin{cases} f(\epsilon) & \text{if } x \in S_\epsilon, \\ 0 & \text{otherwise.} \end{cases}$$

The norm $\|H\|$ of $H \in \mathcal{H}_c(M)$ is defined as

$$\|H\| = \int_0^1 \left(\sup_{x \in M} H(t, x) - \inf_{x \in M} H(t, x) \right) dt.$$

We say that $H, K \in \mathcal{H}_c(M)$ are equivalent, $H \sim K$, if $\varphi_H = \varphi_K$ and the paths $\{\varphi_H^t\}, \{\varphi_K^t\}$, $t \in [0, 1]$, are homotopic with fixed end points. Following [34] we set

$$\mu(F, s) = \inf \{ \|H\| \mid H \sim sF \}, \quad s \geq 0,$$

and define the asymptotic non-minimality $\mu(F)$ of F as

$$\mu(F) = \lim_{s \rightarrow \infty} \frac{\mu(F, s)}{\|F\|}.$$

Then $\mu(F) \in [0, 1]$. Since the support of F is displaceable, it follows from [1, Theorem 3.3.A] or [35, Theorem 8.3.A] that

$$\mu(F) \leq \frac{1}{2},$$

see also [20]. On the other hand, our choice of F guarantees that the Hessian of F at any of its critical points vanishes. If the flow φ_F^s , $s \geq 0$, had no non-constant contractible closed orbit, then sF were “slow” for every $s \geq 0$ in the terminology of [31], and so, according to Theorem 1.4 in [31],

$$\mu(F) = 1,$$

see also [5, 30, 32]. This contradiction shows that the flow φ_F^s has a non-constant contractible closed orbit, and so there exists $\epsilon \in]\epsilon_0 - 2\delta, \epsilon_0 + 2\delta[$ such that $\mathcal{P}^0(S_\epsilon) \neq \emptyset$. Since $\epsilon_0 \in I$ and $\delta > 0$ sufficiently small were arbitrary, Theorem 1.1 follows. $\square$

Proof of Corollary 1.2: Choose $\epsilon > 0$ so small that the local flow ψ_t of the Liouville vector field X defined near S exists for all $t \in]-\epsilon, \epsilon[$. Then ψ_t induces a bijection $\mathcal{P}^0(S) \rightarrow \mathcal{P}^0(\psi_t(S))$, $x \mapsto \psi_t x$, see [17, p. 122]. Corollary 1.2 thus follows from Theorem 1.1. $\square$

Proof of Corollary 1.3: We closely follow [7, Section 12.1]. Let (N, g) and (T^*N, ω_σ) be as in Corollary 1.3. Since σ does not vanish, $\dim N \geq 2$, and so every energy level $E_c = \{H = c^2/2\}$, $c > 0$, is a connected hypersurface. Let $\chi(N)$ be the Euler characteristic of N .

Case 1. $\chi(N) = 0$. We define $d \in [0, \infty[\cup \{\infty\}$ by

$$d = d(g, \sigma) = \sup \{ c \geq 0 \mid H^c \text{ is displaceable in } (T^*N, \omega_\sigma) \}.$$

Notice that

$$d = \sup \{c \geq 0 \mid E_c \text{ is displaceable in } (T^*N, \omega_\sigma)\}.$$

Since $\sigma \neq 0$, the zero section N of T^*N is not Lagrangian, and so a remarkable theorem of Polterovich [33, 22] implies that $d > 0$. We can assume without loss of generality that d is finite.[†] Corollary 1.3 follows from applying Theorem 1.1 to $S = E_{d/2}$ and a thickening

$$\psi: S \times]-d/2, d/2[\rightarrow \bigcup_{0 < c < d} E_c$$

such that $\psi(S \times \{\epsilon\}) = E_{\epsilon+d/2}$.

Case 2. $\chi(N) \neq 0$. In this case the zero section N is not displaceable for topological reasons. We use a stabilization trick used before by Macarini [27]. Let S^1 be the unit circle, and denote canonical coordinates on T^*S^1 by (x, y) . We consider the manifold $T^*(N \times S^1) = T^*N \times T^*S^1$ endowed with the split symplectic form $\omega = \omega_\sigma \oplus \omega_{S^1}$, where $\omega_{S^1} = dy \wedge dx$. Then $N \times S^1$ is not Lagrangian, and $\chi(N \times S^1) = 0$. Let

$$H_1(q, p) = \frac{1}{2}|p|^2, \quad H_2(x, y) = \frac{1}{2}|y|^2, \quad H(q, p, x, y) = \frac{1}{2}|p|^2 + \frac{1}{2}|y|^2$$

be the metric Hamiltonians on T^*N , T^*S^1 and $T^*N \times T^*S^1$. In order to avoid confusion, we denote their energy levels by $E_c(H_1)$, $E_c(H_2)$ and $E_c(H)$. Repeating the argument given in Case 1 for the Hamiltonian system

$$(2) \quad H: (T^*N \times T^*S^1, \omega) \rightarrow \mathbb{R}$$

and

$$d = d(g, \sigma) = \sup \{c \geq 0 \mid H^c \text{ is displaceable in } (T^*N \times T^*S^1, \omega)\}$$

we find that $\mathcal{P}^\circ(E_c(H)) \neq \emptyset$ for a dense set of $c \in]0, d[$. Fix $c \in]0, d[$ such that $\mathcal{P}^\circ(E_c(H)) \neq \emptyset$. Since the Hamiltonian system (2) splits, a contractible closed orbit $x(t)$ on $E_c(H)$ is of the form $(x_1(t), x_2(t))$, where x_1 is a contractible closed orbit on $E_{c_1}(H_1)$ and x_2 is a contractible closed orbit on $E_{c_2}(H_2)$ and $c_1 + c_2 = c$. Since the only contractible orbits of $H_2: T^*S^1 \rightarrow \mathbb{R}$ are the constant orbits on $E_0(H_2)$, we conclude that $c_2 = 0$ and $c_1 = c$, and so $x_1 \in \mathcal{P}^\circ(E_c(H_1))$. The proof of Corollary 1.3 is complete. $\square$

[†]If σ is exact, then $d(g, \sigma)$ is not larger than Mané's critical value and in particular finite, see [7, Section 12.2]. If σ is the area form on the flat 2-torus T^2 , then (T^*T^2, ω_σ) is symplectomorphic to $(T^2, \sigma) \times (\mathbb{R}^2, \omega_0)$, see [4, Section 2.4.B], and so $d(g, \sigma) = \infty$.

Proof of Corollary 1.4: Arguing by contradiction we assume that L and hence a neighbourhood U of L in M is displaceable. We choose a Riemannian metric as in (ii) and denote by $|p|$ the length of a covector $(q, p) \in T^*L$. By Weinstein's Theorem we find $\epsilon > 0$ such that a neighbourhood of L in U can be symplectically identified with $T_{2\epsilon}^*L = \{(q, p) \in T^*L \mid |p| < 2\epsilon\}$. Then $S_\epsilon = \{(q, p) \in T_{2\epsilon}^*L \mid |p| = \epsilon\}$ is a displaceable hypersurface of contact type, and hence, by Corollary 1.2, carries a closed characteristic which is contractible in M . It corresponds to a closed orbit of the geodesic flow on S_ϵ which is contractible in M . According to (i) and (ii) such orbits do not exist, however. $\square$

REFERENCES

- [1] M. Bialy and L. Polterovich. Invariant tori and symplectic topology. *Amer. Math. Soc. Transl.* **171** (1996) 23–33.
- [2] W. Chen. Pseudo-holomorphic curves and the Weinstein conjecture. *Comm. Anal. Geom.* **8** (2000) 115–131.
- [3] K. Cieliebak, V. Ginzburg and E. Kerman. Symplectic homology and periodic orbits near symplectic submanifolds. [math.DG/0210468](#)
- [4] Y. Eliashberg and L. Polterovich. Unknottedness of Lagrangian surfaces in symplectic 4-manifolds. *Internat. Math. Res. Notices* (1993) 295–301.
- [5] M. Entov. K-area, Hofer metric and geometry of conjugacy classes in Lie groups. *Invent. Math.* **146** (2001) 93–141.
- [6] A. Floer, H. Hofer and C. Viterbo. The Weinstein conjecture in $P \times \mathbb{C}^l$. *Math. Z.* **203** (1990) 469–482.
- [7] U. Frauenfelder and F. Schlenk. Hamiltonian dynamics on convex symplectic manifolds. [math.SG/0303282](#).
- [8] V. Ginzburg. On closed trajectories of a charge in a magnetic field. An application of symplectic geometry. Contact and symplectic geometry (Cambridge, 1994), 131–148, *Publ. Newton Inst.* **8**, Cambridge Univ. Press, Cambridge, 1996.
- [9] V. Ginzburg. Hamiltonian dynamical systems without periodic orbits. Northern California Symplectic Geometry Seminar, 35–48, *Amer. Math. Soc. Transl. Ser. 2*, **196**, Amer. Math. Soc., Providence, RI, 1999.
- [10] V. Ginzburg and B. Gürel. Relative Hofer–Zehnder capacity and periodic orbits in twisted cotangent bundles. [math.DG/0301073](#).
- [11] V. Ginzburg and E. Kerman. Periodic orbits of Hamiltonian flows near symplectic extrema. *Pacific J. Math.* **206** (2002) 69–91.
- [12] M. Gromov. Pseudo-holomorphic curves in symplectic manifolds. *Invent. math.* **82** (1985) 307–347.
- [13] H. Hofer and C. Viterbo. The Weinstein conjecture in cotangent bundles and related results. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. IV* **15** (1988) 411–445.
- [14] H. Hofer and C. Viterbo. The Weinstein conjecture in the presence of holomorphic spheres. *Comm. Pure Appl. Math.* **45** (1992) 583–622.

- [15] H. Hofer and E. Zehnder. Periodic solutions on hypersurfaces and a result by C. Viterbo. *Invent. Math.* **90** (1987) 1–9.
- [16] H. Hofer and E. Zehnder. A new capacity for symplectic manifolds. *Analysis, et cetera*, 405–427, Academic Press, Boston, MA, 1990.
- [17] H. Hofer and E. Zehnder. *Symplectic Invariants and Hamiltonian Dynamics*. Birkhäuser, Basel, 1994.
- [18] M.-Y. Jiang. Hofer–Zehnder symplectic capacity for two-dimensional manifolds. *Proc. Roy. Soc. Edinburgh Sect. A* **123** (1993) 945–950.
- [19] E. Kerman. Periodic orbits of Hamiltonian flows near symplectic critical submanifolds. *Internat. Math. Res. Notices* (1999) 953–969.
- [20] F. Lalonde and D. McDuff. Hofer’s L^∞ -geometry: energy and stability of Hamiltonian flows, part I. *Invent. Math.* **122** (1995) 1–33.
- [21] F. Lalonde and L. Polterovich. Symplectic diffeomorphisms as isometries of Hofer’s norm. *Topology* **36** (1997) 711–727.
- [22] F. Laudenbach and J.-C. Sikorav. Hamiltonian disjunction and limits of Lagrangian submanifolds. *Internat. Math. Res. Notices* 1994, 161–168.
- [23] G. Liu and G. Tian. Weinstein conjecture and GW-invariants. *Commun. Contemp. Math.* **2** (2000) 405–459.
- [24] G. Lu. The Weinstein conjecture on some symplectic manifolds containing the holomorphic spheres. *Kyushu J. Math.* **52** (1998) 331–351.
- [25] G. Lu. The Weinstein conjecture in the uniruled manifolds. *Math. Res. Lett.* **7** (2000) 383–387.
- [26] R.-Y. Ma. Symplectic capacity and the Weinstein conjecture in certain cotangent bundles and Stein manifolds. *Nonlinear Differential Equations Appl.* **2** (1995) 341–356.
- [27] L. Macarini. Hofer–Zehnder capacity and Hamiltonian circle actions. [math.SG/0205030](#).
- [28] L. Macarini. Hofer–Zehnder semicapacity of cotangent bundles and symplectic submanifolds. [math.SG/0303230](#).
- [29] L. Macarini and F. Schlenk. A refinement of the Hofer–Zehnder theorem on the existence of closed trajectories near a hypersurface. Preprint 2003.
- [30] D. McDuff. Geometric variants of the Hofer norm. *J. Symplectic Geom.* **1** (2002) 197–252.
- [31] D. McDuff and J. Slimowitz. Hofer–Zehnder capacity and length minimizing Hamiltonian paths. *Geom. Topol.* **5** (2001) 799–830.
- [32] Y.-G. Oh. Chain level Floer theory and Hofer’s geometry of the Hamiltonian diffeomorphism group. [math.SG/0104243](#).
- [33] L. Polterovich. An obstacle to non-Lagrangian intersections. The Floer memorial volume, 575–586, *Progr. Math.* **133**, Birkhäuser, Basel, 1995.
- [34] L. Polterovich. Geometry on the group of Hamiltonian diffeomorphisms. Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998). *Doc. Math.* **1998**, Extra Vol. II, 401–410.
- [35] L. Polterovich. *The geometry of the group of symplectic diffeomorphisms*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2001.
- [36] C. Viterbo. A proof of Weinstein’s conjecture in $\mathbb{R}^{2n}$. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **4** (1987), 337–356.
- [37] C. Viterbo. Exact Lagrange submanifolds, periodic orbits and the cohomology of free loop spaces. *J. Differential Geom.* **47** (1997) 420–468.

- [38] C. Viterbo. Functors and computations in Floer homology with applications. I. *Geom. Funct. Anal.* **9** (1999) 985–1033.
- [39] A. Weinstein. On the hypotheses of Rabinowitz' periodic orbit theorems. *J. Differential Equations* **33** (1979) 353–358.

(U. FRAUENFELDER) DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN

E-mail address: `urs@math.sci.hokudai.ac.jp`

(F. SCHLENK) ETH ZÜRICH, CH-8092 ZÜRICH, SWITZERLAND

E-mail address: `schlenk@math.ethz.ch`